

003951899

WPI Acc No: 84-097443/198416

XRAM Acc No: C84-041481

Antiinflammatory 4-hydroxy-2-thiophene carboxylic acid derivs. - useful intermediates for medicines and pesticides, and as medical drugs

Patent Assignee: MITSUBISHI PETROCHEMICAL CO LTD (MITP ); MITSUBISHI YUKA YAKUHIIN KK (MITP )

Number of Countries: 001 Number of Patents: 001

Patent Family:

Patent No	Kind	Date	Applicat No	Kind	Date	Main IPC	Week
JP 59042375 A		19840308	JP 82153595 A		19820903		198416 B

Local Applications (No Type Date): JP 82153595 A 19820903

Priority Applications (No Type Date): JP 82153595 A 19820903

Patent Details:

Patent	Kind	Lan Pg	Filing Notes	Application	Patent
JP 59042375 A		9			

Abstract (Basic): JP 59042375 A

Cpds. of formula (I), their salts and esters are new, and have antiinflammatory effect: (R1 is H or 1-4C alkyl; R2 is H, 1-6C alkyl, opt. substd. phenyl, benzoyl, or substd. ben oyl).

42.8% inhibition of carrageenin edema is exhibited in rats when (I) is administered in an oral dose of 25 mg/kg.S In an example, 200 mg of 4-hydroxy-5 -phenyl-2-thiophene carboxylic acid ethyl ester (0.76 mmol) was dissolved in 2 ml of 1N NaOH. To this mixt. 0.1 ml of dimethyl sulphate (1.0 mmol) was added dropwise under stirring. The reaction mixt. was heated under reflux for 2 hrs. After cooling, 5 ml of 10% NaOH aq. soln. was added and stirred at room temp. for 1 hr. The mixt. was adjusted at acidic pH with 10% HCl, extracted with ethyl acetate, and conc. When the residue was recrystallised from cyclohexane-toluene (1:1). 106 mg of 4-methoxy-5 -phenyl-2-thiophene -carboxylic acid was obtd.

Title Terms: ANTIINFLAMMATORY; HYDROXY; THIOPHENE; CARBOXYLIC; ACID; DERIVATIVE; USEFUL; INTERMEDIATE; MEDICINE; PEST; MEDICAL; DRUG

Derwent Class: B03; C02

International Patent Class (Additional): C07D-333/32

?

TI- Antiinflammatory 4-hydroxy-2-thiophene carboxylic acid derivs. - useful intermediates for medicines and pesticides, and as medical drugs

AB- J59042375 Cpds. of formula (I), their salts and esters are new, and have antiinflammatory effect: (R1 is H or 1-4C alkyl; R2 is H, 1-6C alkyl, opt. substd. phenyl, benzoyl, or substd. ben<sup>o</sup>yl).

- 42.8% inhibition of carrageenin edema is exhibited in rats when (I) is administered in an oral dose of 25 mg/kg.S In an example, 200 mg of 4-hydroxy-5-phenyl-2-thiophene carboxylic acid ethyl ester (0.76 mmol) was dissolved in 1 ml of 1N NaOH. To this mixt. 0.1 ml of dimethyl sulphate (1.0 mmol) was added dropwise under stirring. The reaction mixt. was heated under reflux for 2 hrs. After cooling, 5 ml of 10% NaOH aq. soln. was added and stirred at room temp. for 1 hr. The mixt. was adjusted at acidic pH with 10% HCl, extracted with ethyl acetate, and conc. When the residue was recrystallised from cyclohexane-toluene (1:1). 106 mg of 4-methoxy-5-phenyl-2-thiophene-carboxylic acid was obtd.

PN- JP59042375 A 840308 DW8416 009pp

PR- JP820153595 820903

PA- (MITP ) MITSUBISHI PETROCHEMICAL CO LTD

- (MITP ) MITSUBISHI YUKA YAKUHI KK

MC- B07-B01 B12-D07 C07-B01 C12-D07

DC- B03 C02

IC- C07D333/32

AN- 84-097443 [25]

